113

Tikrit Journal of Pure Science

Scientific Annual Journal Issued by College of Science / University of Tikrit

Editorial Board

Prof. Dr. Ibrahim Shaban Daoud

Asst. Prof. Dr. Sabbar Abdallah Salih

Asst. Prof. Dr. Ali Taha Ali

Asst. Prof. Dr. Awatif Sabir Jassim

Asst. Prof. Dr. Akeel Hussein Ali

Dr. Fatima Abodi Ali

Chief Editor

Secretary Editor

Member

Member

Member

Member

Consultation Board

Prof. Dr. Salem Hassan Al-Kutubi

Prof. Dr. Awwad Shaban Daoud

Prof.Dr. Qusai Abdalgadir Al-Chalaby

Prof.Dr. Abdulmajeed Aiada Ibraheem

Asst. Prof. Dr. Liftaa Selman Kadim

Dr. Muaiad Gasid Jalhoom

Dr. Fayadh Mohamed Abid

College of Science/ University of Kufa

College of Science/ University of Tikrit

College of Science/ University of Mosul

College of Education/ University of Tikrit

College of Science/ University of Tikrit

Ibn Sina Company/ Baghdad

College of Science/ University of Tikrit

Administrative Supervision

Saad Mohamed Ahmed

All Correspondence should be addressed to:

Secretary Editor, Tikrit Journal of Pure Science, College of Science, University of Tikrit, P.O. Box. 42, Tikrit, Iraq

E-mail tik jsci@yahoo.co.uk

Table of Content

No.	Title	Page			
21	Hypoglycemic and Hypolipidemic Effects of Grapefruit Juice in Diabetic Rats Wafaa Mohammad Ali Hamed ¹ , Khadija Younis Abid ² , Safaa Abd-Ul aziz Al-Amin ³ 'Department of Clinical Pharmacy, College of Pharmacy, University of Mosul, Iraq 'Department of Pharmaceutical Sciences, College of Pharmacy, University of Mosul, Iraq 3 Department of Chemistry, College of Science, University of Mosul, Iraq				
22	Effects of Some physical and chemical factors of Lower Zab water on Tigris River Reidh A. Abdul Jabar ¹ , Ali. A-Z. Al-Lami ² , Rushdi S. Abdul Khader ³ and Aseel G. Rahdi Department of Biology, College of Science, University of Tikrit, TiKrit, Iraq Ministry of Environment, Iraq Department of Biology, College of Education, University of Tikrit, TiKrit, Iraq	132 – 142			
23	Comparative Differential Thermal Analysis of Epoxy Resin blends with Amic Acid- Copper Complexes Akram A. Mohamed and Ibtehaj Z. Al-Halim Department of Chemistry, College of Science, University of Mosul, Mosul, Iraq	143 – 146			
24	Synthesis and antibacterial activity of some new 1,3,4-oxadiazoles and 1,3,4-thiadiazoles K.M. Daoud ¹ , A.W. Al-Obaydi ¹ and M.J. Mohammed ² ¹ Department of Chemistry, College of Education, University of Mosul, Mosul, Iraq ² Department of Biology, College of Education, University of Mosul, Mosul, Iraq	147 – 151			
25	Synthesis and Characterization of Heterobinuclear Complexes of Amoxicillin Khalaf I. Khallow ¹ and Nuhad A. Al-Omari ² Department of Chemistry, College of Education, University of Mosul, Mosul, Iraq Department of Pharmaceutical Science, College of Pharmacy, University of Mosul, Mosul, Iraq	152 – 155			
26	Synthesis and Characterization of Schiff Bases (derived from p-substituted aniline) and Their Complexes With Co(II), Ni(II) and Cu(II) Akram A. Mohammed Department of Chemistry, College of Education, University of Mosul, Mosul, Iraq	156 – 158			
27	Synthesis and Characterization of Some Schiff Bases Complexes With Co(II) and Ni(II) Najla H. Taher¹ and Akram A. Mohammed² Department of Chemistry, College of Education for Girls, University of Mosul, Mosul, Iraq Department of Chemistry, College of Education, University of Mosul, Mosul, Iraq	159 – 161			
28	Synthesis of some aryl amino acetyl -2-amino-4-nitrobenzothiazole Derivatives Khalid A. Al-Badraani Department of Sciences Food & Biotechnology, College of Agriculture, University of Tikrit, Tikrit, Iraq	162 – 165			
29	Synthesis of some Phthalazine from Hydrazone of Amino Acids H.A. Basheer ¹ and N.S. Ezzat ² Department of Science, College of Basic Education, University of Mosul, Mosul, Iraq Department of Chemistry, College of Education, University of Mosul, Mosul, Iraq	166 – 170			
30	Trivalent Metal Complexes of 1,3,4-Thiadiazole Derivatives Khalil K. Abid Department of Chemistry, College of Science, University of Al-Mustanserya, Baghdad, Iraq	171 – 175			
31	Utilizing Calcium Carbide for Aqueous Ethanol Dehydration Azzam A. Ismael ¹ , Rafie R. Mohammad ² and Huda A. Younis ² 'College of Technique, Mosul, Iraq ² Technical Institue, Mosul Iraq	176 – 178			
32	Effect of Curvature on Stability of Compressible Fluid Flow in circular Pipe Section Wafaa Muhi Aldeen Department of Mathematics, College of Education, University of Tikrit, Tikrit, Iraq	179 – 181			
33	Weakly regular and pre weakly regular sets in Intuitionistic fuzzy special topological spaces Fatima Mahmmod Mahmmad Department of mathematic, College of Education, University of Tikrit, Tikrit, Iraq	182 – 184			
34	Generalized SF-Rings Raida D. Mahmood ¹ and Abdullah M. Abdul-Jabbar ² Department of Mathematics, College of Computer and Mathematical Science, University of Mosul, Mosul, Iraq Department of Mathematics, College of Science, University of Salahaddin-Erbil, Kurdistan Region, Iraq	185 – 188			
35	MCD-Domain of type A+x B[x] and A+xB[[x]] Mahera Rabee Q. Department of Mathmatics, College of Education, University of Tikrit, Tikrit, Iraq	189 – 191			
36	Analysis of the Longitudinal Profiles of the Doluge Valley in Nineve Governorate using GPS Ziaad R. Elias College of Education, University of Salah All-Deen, Erbil, Iraa	192 – 197			

INSTRUCTIONS TO AUTHORS

Aims and Scope:

The Tikrit Journal of Pure Science (Tikrit, J.Sci) is a biannual refereed periodical devoted to the publication of articles that conform to its scope and editorial standards. The scope encompasses various disciplines of experimental and theoretical original studies in pure and applied sciences.

The Journal publishes original research articles, review articles, short communications, case studies, and letters to the editor. Abstracts of meetings and symposia will also be considered for publication. All submitted papers are subjects to an extensive peer review by a panel of referees.

Submission

Contributions should be sent to the Executive Editor: Chief Editor

College University of Tikrit-Iraq, E.mail: tik jsci@vahoo.co.uk

The original manuscript and two copies should be submitted. The manuscript should be written in a good English or Arabic language. Submitted papers will be referees and, if necessary, authors may be asked to revise their manuscripts.

\$ \$\$ 我我想得我妈妈妈我妈妈我想得我想到我我想到我的事情的我想到我我想到我的感觉的感觉,因为我们是必要感觉

By submitting a paper for publication in Tikrit Journal of pure science the authors state explicitly that the paper has not been published nor has been submitted totally or partially for publication elsewhere, the author has no right to withdraw his article while it is under processing unles he presents a written paper to do so and has to pay for refereeing expenses.

Time Schedule

The main author should receive acknowledgement of receipt of the manuscript and within few months, he should have a notification of acceptance or rejection or a request for revising.

Reprints:

One reprint will be supplied to the principal author free of charge.

Copyright Transfer:

Upon publishing an article, its copyright is transferred automatically to the publisher without Further notification to the author.

Manuscript preparation

In preparation of the manuscript, the following instructions should be strictly. Followed:

1-Manuscripet: the original manuscript should be typed on one side of A4 paper (210mm.X.297mm). Double-spaced with a 35mm. Margin Alleges should be numbered Authors are requested to use SI. Units as the primary listing and to include., in parentheses, other conventional units only when their Listing is extremely necessary to improve immediate understanding of the article contents. Authors are asked after acceptance of their paper, to send the manuscript on a disc indicating the software used.

2-Layout: Authors are requested to adhere to the following order:

Title authors(s), Affiliations, Abstract, Keywords, Main text (1-Introduction 2-Experimental 3-results and Discussion, 4-Aknowledgments, 5-Appendices, 6-References, Figures, Legends and Tables.

- 3.Length: The total length of the paper should not exceed 10 printed pages of the Journal.
- 4. Title: The title should be concise and brief.
- 5. Abstract: The abstract should not exceed 200 words. It should contain the key words of the papers as well as the essence of the results achieved.
- **6.** Figures: Each Figure must be referred to in the text, and have a caption below it. The captions should be indicated in the margin axes of figures must be labeled properly. The drawings for the figures must be copied by scanner and inserted in suitable location after the paragraph that refers to the figure directly. The figures should be identified by the name and figure number.

Tables:

Tables should be inserted in a suitable location after the paragraph that refers to the table directly and each table should have a number and headed by a tittle.

References:

References to other works should be consecutively numbered in which they are cited in the text identified by Arabic numerals in big parentheses [], and should be listed by number on a separate sheet at the end of the paper. Examples of correct reference forms are given below.

An article/ paper in a Journal / Peroiodical / Serial

S.A. Al-jibori, O.H. Amin, T.A.K. Al-Allaf, R.Davis.

A Book

W.R. Adams (1977) " Developing reading versatility" ,2nd Ed. New York Holt, Rinehart and Winston.

A Chapter in a Book

K. Koerner (1994) Historiography of linguistics in:the encyclopedia of languages and linguistics (Eds. Asher Rand Simpson J) Oxford: Pergamon press Vol. 3,P.1572.

A Thesis

J.N. Edwards (1990) Ph.D. thesis, University of Bristol, U.K.

A Patent

Y. Funatsukuri (1996) (Sanyo Electric Co.) projection type colors liquid crystal device Japanese Patent JP 08.5, 1475

Internet www Address

B. Hodsdon (1998) A different Kind of Cinema the silent Years. National library of Australia at, URL: http://www.einemedia.net/NLA.csilent.html

Effects of Some physical and chemical factors of Lower Zab water on Tigris River

Reidh A. Abdul Jabar¹, Ali. A-Z. Al-Lami², Rushdi S. Abdul Khader³ and Aseel G. Rahdi

Department of Biology, College of Science, University of Tikrit, TiKrit, Iraq

² Ministry of Environment, Iraq

³ Department of Biology, College of Education, University of Tikrit, TiKrit, Iraq

Abstract:

Four stations were selected for water sample collections, to study the effects of some water characteristics of Lower Zab tributary on the Tigris river, during the period from November 2001 to October 2002.

Air and water temperature were (10-38)°C and (10-28)°C respectively. Salinity values ranged (0.25-0.33)%_o. Turbidity varied from (22) NTU at station 2 to (34.4) NTU at station 3. pH values ranged (7.5 - 8.6). Dissolved Oxygen values were higher in Lower Zab water compared to that of Tigris River. BOD were high at summer and low during winter.

Total hardness ,calcium and Alkalinity were (222-306) mg/L, (133-165) mg/L and (134-165) mg/L respectively. Nutrition values of the water samples were low, $(0.4\mu g/L)$ for nitrite, $(0.27\mu g/L)$ for phosphate, and (2.1

Introduction:

Surface water in Iraq forms an area of (2400km²). This forms 5% of the total Iraqi lands, which is distributed in various water systems [1].

Hydrological characteristics of Iraqi lands affected by the two rivers, Tigris and Euphrates and their related branches [2]. Lower Zab tributary forms one of the major branches of Tigris River, with a length extends 400 km and over an area of (22250 km²) from its origin in Iran. This area is mainly located in Zakaros mountain in Iran. In Iraq, its length extends about 175km and above 200m wide [3]. Lower Zab consists of a gravel base with a mean water discharge of (233-245)m³/sec [4].

Water characteristics of Rafidain valley like many other water areas of the world is affected by the climate, nature, soil, inner water and its bioactivities. In addition, Tigris river limnology is generally affected by it's tributaries [2].

Many studies conducted on the Iraqi surface water, but were all concentrated on limited area [5, 6, 7, 8]. Therefore, the present study aims to know the effects of some physical and chemical factors of Lower Zab water on Tigris River water.

mg/L) for silica at station 4 after the confluence. While these values were higher at station 3 before the confluence $(0.6\mu g/L)$, $(0.4\mu g/L)$ and (2.9 mg/L)respectively. Chloride ions values ranged (33.5-38.6) mg/L .Discussion of the results of this study clarify that Lower Zab effects the environment of the Tigris river at station 4 through the values of the turbidity, total hardness and negative iones which is located in the river after the confluence, in addition to the decrease of nutritive compared to station 3 in the river. This is certainly due to the favorable conditions at station 4, which cause blooming and diversity of aquatic organisms. This was obvious from the BOD5 values at station 4. The high BOD5 values was proportional to organic pollutants.

Materials and Methods:

Four stations were selected in Tigris river (stations 3,4) and in the Lower Zab tributary (stations 1,2) (Figure -1). Monthly water samples were collected from November 2001 to October 2002.

Air and water temperatures were measured using mercury thermometer. A conductivity meter (YSI-33cc) was used to measure water conductivity (μ semins/cm). Salinity was calculated using the following equation [9]:

Salinity (gm/L) = 0.00064 X conductivity. Turbidity (NTU) was measured using the turbidity meter (HACH-16800cc). Water pH was measured using a pH meter (pH-EC. TDC meter /HI-9811). Winkler method [10] was used to measure dissolved oxygen (mg/L) and Biochemical Oxygen Demand (BOD). The method described in [11] was used to measure alkalinity, total hardness and calcium ions. Chloride, Nitrate, active phosphate and Silica were measured using the methods in standard methods for the examination of water method for Sea water analysis [10, 12].




Fig. 1: Study Area; Lower Zab (stations 1, 2) & Tigris River (stations 3, 4)

Results & Discussion:

Air and water temperatures were close. In that, both air and water temperatures increased during summer and early autumn, and decreased in winter (Table-1; Figure 2). Salinity values varied at the different stations. Low values recorded at station 4 in Tigris river (0.25 gm/L), and the maximum values recorded in Lower Zab at station 2 (0.33 gm/L) (Table-1; Figure 3-a). Average turbidity values in Tigris River at station 4 and 3 were (26.5 NTU) and (34.4 NTU) respectively. While turbidity at stations 1 and 2 were in between the previous values (Table-1; Figure 3-b). Turbidity in the rivers are at the almost high during increase water level and high rapid flow, Which causes increase water suspended matter and mixing of water during rain periods(Hynes 1974). In a study on Tigris River [2], turbidity values were related to the suspended solid substances, such as mud and silts which increased during the periods of rains and high discharge.

pH values in the present study inclined towards weak alkalinity (7.5-8.6) (Table-1: Figure 5-a). This coincides with results of previous studies [13, 14, 15], which pointed towards weak alkalinity of Iraqi water.

Dissolved oxygen values did not fall below 5.2 mg/L. High values recorded in the Lower Zab compared to Tigris River (Table-1: Figure 4-a). This was related to the fast flow and good mixing [14, 15, 16]. Biochemical Oxygen Demand (BOD₅) values were high during summer (Table-1; Figure 4-b). The real cause is the presence of organic biodegradable matter that are consumed by bacteria. Dissolved oxygen, also decreases with an increase in temperature [17].

Total hardness recorded high values at station 3 in Tigris River, compared with station 1 in Lower Zab (Table-1; Figure 6-a). AL- Lami [2] founded close values, and this was attributed as follows; Tigris river water is considered hard water, and the real causes of hardness is the nature of catchments area through which the Tigris river flows.

Also the polluted loads discharges add to this hardness. Low calcium ions values (133 mg/L) recorded at station 4 in Tigris River, and this could be due to the consumption of the ions by living organism bloomed at this station [3]. While, higher calcium ion values (165 mg/L) recorded at station 2 in Lower Zab. This is due to the surrounding soil nature (Table-1; Figure 6-b).

Table-1 and Figure 5-b, shows alkalinity values between 52 and 208 mg/L. These values were within the natural range [10]. Because most water alkalinities in Tigris River are due to bicarbonate ions. This was pointed out in many previous studies [8, 13].

Mean nitrite values recorded in the present study at all stations were close and low. A decreased value of nitrite below sensation level was recorded in station 3 in Tigris River (Table-1; Figure 7-b). Nitrite values in the unpolluted good ventilated waters are low [2]. This is true because of the values of dissolved oxygen at the same station (6.5- 9.7 mg/L).

Phosphate values at stations 3 and 4 were (0.4 $\mu g/L$) and (0.27 $\mu g/L$) respectively in Tigris River (Table-1; Figure 8-a). This is because of dilution effect after the confluence of Lower Zab into Tigris River. Also due to the consumption by living aquatic organisms. Plant nutritives considered to be very important in increasing primary productivity, which forms the principal base of food chain in aquatic environment [18].

The maximum silica value (5.7 mg/L) recorded at station 1 in Lower Zab, and the lower values recorded at station 4 in Tigris River (2.1 mg/L) (Tabe-1; Figure 8-b). Silica found naturally in water in concentrations ranged (1-10 mg/L). Silica concentration ranged (0.5-0.8) mg/L required for diatoms growth [19].

The upper chloride values recorded at all stations did not exceeded 200 mg/L ; the upper chloride ions concentration allowed and prefered in drinking water recommended by WHO (1971). And the mean values recorded in the present study were close to the preferable concentration of 25 mg/L of drinking water .The low chloride value recorded was 7.5 mg/L .This was lower than the values recorded by Saleh [20] and Al-Doori [21] in Tigris river /Salahudeen province. The upper chloride value was (38.6 mg/L), this was also lower than the values recorded by the previous two authors .The mean chloride values were close to the preferable level (25 mg/L) allowed in the international characterization and the characterization of Iraqi waters No. 25 in 1967. Chlorides are among the important negative ions in natural waters and which give the water its salty test when it binds with sodium ions and change with concentration. One of the most special characters of Iraqi waters are the dominancy of Ca, HCO₃, Cl, SO₄ ions .This emphasize the mineral contents of the rocks in the river basin, which was also stated by Al-Lami and Al-Aubaidi [14].

Table 1: Range, Mean and standard deviations of the physical and chemical measurements for the surface water at the different stations during 2001-2002.

Station	Lower Zab		Tigris river	
Factors	1	2	3	4
Air Temperature (°C)	12 - 36	12 -34	13.5-38	10-31
Water Temperature (°C)	11 - 27	10 – 27	10-28	10-24
Salinity (%0)	0.19 - 0.46 $(0.31) \pm 0.1$	0.2 - 0.59 $(0.33) \pm 0.1$	0.22-0.46 (0.28) =0.1	0.16-0.35 $(0.25) \pm 0.04$
Turbidity (NTU)	0.73 - 111 $(27.1) \pm 30$	ND - 109 (22) = 30.7	0.24-185 (34.4) ±54.8	2.57-72 (26.5) ±23
рН	$7.9 - 8.3$ $(8.12) \pm 0.1$	$7.7 - 8.4$ $(8.1) \pm 0.1$	7.8-8.3 (8.1) ±0.1	$7.5-8.6$ $(8.1) \pm 0.3$
Dissolved Oxygen (mg/l)	5.2 - 10 $(7.9) \pm 1.2$	6.3 - 10.4 $(7.9) = 1.2$	6.5-9.7 (7.6) =0.9	6.1-10 $(7.7) \pm 1.1$
BOD (mg/l)	0.4 - 4.6 (2.5) = 1.6	0.7 - 4.6 $(2.3) = 1.5$	0.7-3.8 (2) ± 1.1	1.3-4.5 (2.7) ±1.2
Total Hardness (mg/l)	$65 - 540$ $(281) \pm 114$	175-510 (293) ±80	120-635 (306) =135	110-475 (222) ±94
Calcium (mg/l)	50 - 260 (154) ± 68	60-280 (165) ±64	50-245 (141) =54	50-225 (133) ±56
Alkalinity (mg/l)	$104 - 208$ $(152) \pm 0.6$	52 - 208 (135) ± 49	104 – 208 (165) = 41.6	$ \begin{array}{r} 104 - 208 \\ (134) \pm 33 \end{array} $
Nitrite (μg/l)	0.016 - 1.534 $(0.7) \pm 0.6$	0.008-2.6 (0.5) = 0.7	N.D-2.04 (0.6) =0.8	0.07-2.12 $(0.4) \pm 0.5$
Phosphate (µg/l)	$0.012 - 1.23$ $(0.33) \pm 0.3$	0.03-1.24 $(0.33) \pm 0.4$	0.06-1.5 (0.4) =0.3	0.029-1.368 $(0.27)\pm0.3$
Silica (mg/l)	1.98 -18.5 (5.7) ± 4.9	0.06-6.6 (3.8) ±1.5	0.2-10.9 (92.9) = 3.2	$0.06-3.97$ $(2.1)\pm1.1$
Chloride (mg/l)	23.1 - 45.3 $(33.8) \pm 7.5$	26.6 - 65.7 (38.6) = 11.8	22.8 - 52.4 $(34) = 8$	$26.3 - 58.6$ $(33.5) \pm 8.5$

N.D = Not detected.




Fig. (2): Air & Water Temperature at the study area from Nov. 2001 to Oct. 2002




Fig. (3-a&b): Salinity & Turbidity at the study area from Nov. 2001 to Oct. 2002




Fig. (4-a&b): PH &Dissolved Oxygen at the study area from Nov. 2001 to Oct. 2002




Fig. (5-a&b): BOD& Total Hardness at the study area from Nov. 2001 to Oct. 2002




Fig. (6-a&b): Calcium & Alkalinity at the study area from Nov. 2001 to Oct. 2002




Fig. (7-a&b): Nitrite & Phosphate at the study area from Nov. 2001 to Oct. 2002




Fig. (7-a&b): Silica & Chloride at the study area from Nov. 2001 to Oct. 2002

المصادر:

- السعدي، حسين علي (1994). البيئة المائية في العراق و مصادر تلوثيا.
 وقائع مؤتمر البحث العلمي ودوره في حصية البيئة من مخاطر التلوث.
 تحرير الدكتور حسين علي السعدي ، دمشق 26-88/9/89 اتحاد مجالس البحث العلمي العربية الامائة العامة ، بغداد 69-88.
- اللامي، على عبد الزهرة (1998). التأثيرات البيئية لذراع الثرثار عنى نير دجلة قبل دخوله مدينة بغداد. رسالة دكتوراه، كلية العلوم، الجامعة المستنصرية .123ص.
- اللامي وجماعته (مقبول للنشر) (2005). اتأثيرات البيئية لرافد الزاب الأسفل في تتوع الافقريات القاع في نهر عجلة. مجلة كلية العلوم جامعة تكريت.
- 4. Sabri; A.W., Kassim; T.I and Shawaket; S.F. (1999). The effect of Lesser Zab river on the Tigris river. IAEC Publ. Rep. No. 90:4/6180.
- 5. الأمارة، فارس جاسم محمد و عليوي ، يسرى جعفر و موسى ، فتن صدام(2001). التغيرات الشهرية في مستويات الاملاح المغنية والكلوروفيل أ في مياه شط العرب. مجلة وادي الرافدين لعلوم البحار. جامعة البصرة. 16(1): 347-357.
- 6. حسن، فكرة مجيد (1988). دراسة بيئية وفسلجية ونوعية للهائمات النباتية في هور الحمار. رسالة ماجستير جامعة البصرة 136 ص.
- حسين، صادق على (2001). مصادر التلوث العضوي في المياه الداخلية العراقية و امكانية السيطرة عليها و اعادة استخدامها. مجلة و ادي الرافدين لعلوم البحار، 1(1): 489-505.

after crossing Baghdad. J. Coll. Educ., women, Univ., Baghdad, 5(1):43-50.

16. اللامي، على عبد الزهرة و مفتن، فاطمة شغيت ونشأت، مهند رمزي (2001). لافقريات القاع في ذراع الثرثار ونهر دجلة. مجلة ابن الهيثم للعلوم الصرفة و التطبيقية، 14(3):87-98.

17. Laws, E.A. Aquatic Pollution . Awileg. Inter Science publication . U.S.A.:482 (1981).

18. السعدي، حسين علي و اللامي، على عبد الزهرة و قاسم، ثائر إبراهيم (2000). العوامل البيئية وعلاقتها بالاستزراع السمكي في بحيرة القادسية. مجلة كلية التربية للبنات، جامعة بغداد. 11(2):35-45.

19. Reid, G.K.(1961). Ecology of inland water and estuaries. Reihod publ. corp., New York.

20. صالح، موفق انهاب(2000). دراسة لمنولوجية على نهر دجلة ضمن محافظة صلاح الدين. رسالة ماجستير، كلية التربية للبنات – جامعة تكريت.

21. الدوري، نهاد عبد محمد(2000). تأثير الملوثات الصناعية والسكنية على نهر دجلة ضمن محافظة صلاح الدين. رسالة ماجستير، كلية التربية – جامعة تكريت.

8. Al- Saadi. H. A., Hadi, R. A. M. & AL-Lami, A.A.(1989). Limnological studies on some marsh areas in southern Iraq. Limnological, 20(2).

9.Richards , L.A.(1954). Dignosis and improvement of Saline and AL-Kalisoils – United States Department of Agriculture , Hand book No. 60. USA. Washington, D.C.160p.

 APHA (1985)., AWWA, WPCE. Standard methods for the examination of Water and Waste Water. 14th. ed.

11. Lind, O.T. (1979). Hand Book of common methods in Limnology. C.V. Mosby, St. Lonis..

 Parsons; T.R., Maita; y. & Lalli; C.M. (1984).
 Amanual of chemical and biological methods for seawater analysis, Pergamon press, oxford.

13. الربيعي، ميادة عبد الحسين جعفر (1997). دراسة بيئية عن نهر العظيم وتأثيره على نهر دجلة. رسالة ماجسير، جامعة بغداد. ص105.

14. اللامي، على عبد الزهرة و العبيدي، خنساء حميد (1996). دراسة بعض الخصائص الفيزيائية والكيميائية خزان الثرثار - العراق. مجلة كلية التربية للبنات ، 7(2):2 - 28.

15. Maulood; B.K., AL- Azzawi; M.N. & Saadall; H.A. (1994). An ecological study on the Tigris river pre and

تأثير بعض العوامل الفيزيائية والكيميائية لمياه الزاب الأسفل في نهر دجلة رياض عباس عبد الجبارا و على عبد الزهرة اللامي2 و رشدي صباح عبد القادر3 و أسيل غالى راضى

أقسم علوم الحياة، كلية العلوم، جامعة تكريت، تكريت، جمهورية العراق وزارة البيئة، جمهورية العراق

· قسم علوم الحياة، كلية التربية، جامعة تكريت، تكريت، جمهورية العراق

الملخص:

اختيرت أربعة محطات لجمع عيدت المياه لدراسة تأثيرات بعض خصائص مياه راقد الزاب الأسفل عنى مياه نير دجلة من تشرين الثاني 2001 ولغاية تشرين الأول 2002 قراوحت درجات حرارة الهواء والماء (10-38)° م و(10-28)° م على التوالي. في حين كانت قيم المفوحة بين (2.0-0.33) جزء بالأف. كانت كدرة المياه(22 وحدة كدرة نقالية) في المحطة 3 و(3.4.4 وحدة كدرة نقالية) في المحطة 3 و(3.4.4 وحدة كدرة نقالية) في المحطة 3 في المحطة 3 براوحت درجة الأس الهيدروجيني في جميع المحطات بين(7.5-8.6). قيم الأوكسجين المذاب كانت عالية في ميد الزاب الأسفل مقارنة بمياه نير نجلة ، في حين كانت قيم المتصب الحيوي للأوكسجين عالية خلال الصيف ومنخفضة خلال الشتاء . سجنت العسرة الكلية وايونات الكالسيوم والقاعدية قيماً تراوحت بين (222-306)منع/تر (133-165) ملغم/لتر على التوالي .

قيم المغذيات لعينات المياه كانت واطنة لحد ما، وبلغت (0.4مايكروغرام / نتر) للنترنيت و(0.27 مايكرو غراد/لتر) للفوسفات و (2.1 ملغم/لتر)

للسليكا في المحطة الرابعة بعد مصب الرافد في نهر دجلة. بينما كانت قيم المغذيات أعلاه في المحطة الثالثة في نهر دجلة مرتفعة، حيث سجلت (0.6 مايكرو غرام/ لتر) و (2.9 ملغم/لتر) على التوالى .

أما قيم الكلوريد فقد تراوحت (33.5-38.6) ملغم/ لتر. نوقشت النتائج على أساس التغيرات البيئية في مياه المحطات او 2و 3 و بالتالي تأثير ها على مياه المحطة الرابعة.وقد تبين من انتائج إن الزاب الأسفل يؤثر على بيئة نهر دجلة في المحطة الرابعة من خلال قيم العكورة، والعسرة الكلية والايونات السالبة والواقعة في نهر دجلة بعد نقطة اتصاله بالنهر، إضافة إلى انخفاض قيم المعذيات مقارنة بالمحطة الثالثة. وهذا بالتأكيد يعود إلى الظروف الملائمة في المحطة الرابعة والتي تسبب الازدهار والتنوع الحياتي للأحياء المائية. وهذا كان واضحاً من قيم المحطة الرابعة العالية للمتطلب الحياتي للأوكسجين في المحطة الرابعة. القيم العالية للمتطلب الحياتي للأوكسجين في المحطة الرابعة. القيم العالية للمتطلب الحياتي